

NEO-OXY (neomycin sulfate, oxytetracycline) Veterinary Feed Directive for use in Calves, Beef and Non-lactating Dairy Cattle

Client: _____ Veterinarian: _____
Business or Home Address: _____
Address: _____
Phone #: _____ Phone #: _____

Approximate number of animals to be treated: _____

Location of animals: _____

Special Instructions and/or other animal identifications:

Indication, Drug Level in Medicated Feed, and Duration of Use (check one):

- A)** For Calves (up to 250 lbs.): For treatment of bacterial enteritis caused by *Escherichia coli* susceptible to oxytetracycline; treatment and control of colibacillosis (bacterial enteritis) caused by *Escherichia coli* susceptible to neomycin.
Drug level (in milk replacers or starter feeds): _____ g/ton in order to provide 10 mg / lb body weight / day
Duration of use: _____ days (specify number of days from 7 to 14 days)
- B)** For Growing Cattle (over 400 lbs.): For the reduction of the incidence of liver abscesses.
Drug level: _____ g/ton in order to provide 75 mg/head/day
Duration of use: _____ days (specify number of days)
- C)** For Cattle: For prevention and treatment of the early stages of shipping fever complex
Drug level: _____ g/ton in order to provide 0.5 – 2.0 g/head/day
Duration of use: _____ days (specify number of days from 3 to 5 days before and after arrival in feedlot)
- D)** For Calves, Beef, and Non-lactating Dairy Cattle: For treatment of bacterial enteritis caused by *Escherichia coli* and bacterial pneumonia (shipping fever complex) caused by *Pasteurella multocida* susceptible to oxytetracycline; treatment and control of colibacillosis (bacterial enteritis) caused by *Escherichia coli* susceptible to neomycin.
Drug level (in feed or milk replacer): _____ g/ton in order to provide 10 mg/lb body weight / day
Duration of use: _____ days (specify number of days from 7 to 14 days)

Caution: Use of feed containing this Veterinary Feed Directive (VFD) drug in a manner other than as directed on the labeling (extra-label use) is not permitted.

For use in Dry Feeds Only. Not for Use in Liquid Medicated Feeds.

Residue Warnings:

Indications A and D: Withdraw 5 days before slaughter. Use of more than one product containing neomycin or failure to follow withdrawal times may result in illegal drug residues. A withdrawal period has not been established for use in preruminating calves. Do not use in calves to be processed for veal. A milk discard time has not been established for use in lactating dairy cattle. Do not use in female dairy cattle 20 months of age or older.

Indication B: Zero-day withdrawal period. Use of more than one product containing neomycin or failure to follow withdrawal times may result in illegal drug residues.

Indication C: Zero-day withdrawal period. Use of more than one product containing neomycin or failure to follow withdrawal times may result in illegal drug residues. A withdrawal period has not been established for use in preruminating calves. Do not use in calves to be processed for veal. A milk discard time has not been established for use in lactating dairy cattle. Do not use in female dairy cattle 20 months of age or older.

Combination Use:

This VFD only authorizes the use of the VFD drug(s) cited in this order and is not intended to authorize the use of such drug(s) in combination with any other animal drugs.

VFD Issuance Date: _____

VFD Expiration Date: _____

Month/Day/Year
(Not to exceed 6 months from issuance date)

Veterinarian's signature: _____

Original – Veterinarian

Copy – Supplier

Copy – Client